

Supplementary data for article:

Rakić, A. A.; Vukomanović, M.; Trifunović, S.; Travas-Sejdic, J.; Chaudhary, O. J.; Horský, J.; Ćirić-Marjanović, G. Solvent Effects on Dopant-Free PH-Falling Polymerization of Aniline. *Synthetic Metals* **2015**, *209*, 279–296.

<https://doi.org/10.1016/j.synthmet.2015.07.031>

Supporting Information

Solvent effects on dopant-free pH-falling polymerization of aniline

Aleksandra A. Rakić^a, Marija Vukomanović^b, Snežana Trifunović^c, Jadranka Travas-Sejdic^d,
Omer Javed Chaudhary^d, Jiří Horský^e, Gordana Ćirić-Marjanović^{a,*}

^a*Faculty of Physical Chemistry, University of Belgrade, Studentski trg 12-16, 11158,
Belgrade, Serbia*

^b*Jožef Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia*

^c*Faculty of Chemistry, University of Belgrade, Studentski Trg 12-16, 11158 Belgrade,
Serbia*

^d*Polymer Electronics Research Centre, School of Chemical Sciences, University of Auckland, 23
Symonds Street, Auckland, New Zealand and MacDiarmid Institute for Advanced Materials and
Nanotechnology, Victoria University of Wellington, P.O. Box 600, Wellington 6140, New Zealand*

^e*Institute of Macromolecular Chemistry, Academy of Sciences of the Czech Republic,
162 06 Prague 6, Czech Republic*

*Corresponding author. Tel. +381113336633; Fax: +381112187133.

E-mail address: gordana@ffh.bg.ac.rs (G. Ćirić-Marjanović)




Figure S1. Solventless LDI mass spectra of (a) PANI-ACN-1.5 and (b) PANI-ACN-24. The insets show enlarged parts of the mass spectra.




Figure S2. Solventless LDI mass spectra of (a) PANI-DMSO-1.5 and (b) PANI-DMSO-24.




Figure S3. Solventless LDI mass spectra of (a) PANI-EtOH-1.5 and (b) PANI-EtOH-24. The insets show enlarged parts of the mass spectra.




Figure S4. Solventless LDI mass spectrum of PANI-Water. The inset shows enlarged part of the mass spectrum.